

DOI: [https://dx.doi.org/10.21123/bsj.2020.17.1\(Suppl.\).0342](https://dx.doi.org/10.21123/bsj.2020.17.1(Suppl.).0342)

Solving Linear Volterra – Fredholm Integral Equation of the Second Type Using Linear Programming Method

Iman A. Dhari

Najwa R. Mustafa

*Muna M. Mustafa**

Received 22/5/2019, Accepted 16/9/2019, Published 18/3/2020



This work is licensed under a [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/).

Abstract:

In this paper, a new technique is offered for solving three types of linear integral equations of the 2nd kind including Volterra-Fredholm integral equations (LVFIE) (as a general case), Volterra integral equations (LVIE) and Fredholm integral equations (LFIE) (as special cases). The new technique depends on approximating the solution to a polynomial of degree $(m - 1)$ and therefore reducing the problem to a linear programming problem (LPP), which will be solved to find the approximate solution of LVFIE. Moreover, quadrature methods including trapezoidal rule (TR), Simpson 1/3 rule (SR), Boole rule (BR), and Romberg integration formula (RI) are used to approximate the integrals that exist in LVFIE. Also, a comparison between those methods is produced. Finally, for more explanation, an algorithm is proposed and applied for testing examples to illustrate the effectiveness of the new technique.

Keywords: Linear Volterra-Fredholm integral equation of the 2nd kind (LVFIE), Linear programming problem (LPP), Quadrature methods.

Introduction:

Integral equations occur naturally in many fields of mechanics and mathematical physics. They also arise as representation formulas for the solutions of differential equations (1).

From the last few years, there has been interest to use the linear and nonlinear programming methods to find a numerical or approximate solution for integral equations. AL-Nasir in 1999 used the linear programming method to find numerical solution of Volterra integral equations of the 2nd kind (2). Also, Saed in 1999 used linear programming method to find numerical solution of Fredholm integral equations of the 1st kind (3). While Kalwi in 1999 used the same procedure to find numerical solution for Fredholm integral equations of the 2nd kind (4). Shua'a in 2005 calculated numerical solution for Volterra integral equation using linear programming problem (5). Kamyad et al. in 2010 proposed a new approach for solving linear and nonlinear Volterra integral equations of the 1st and 2nd kinds first by defining a new problem in calculus of variations, which is equivalent to this kind of problem, then by using the

Department of Mathematics, College of Science for Women, University of Baghdad, Baghdad-Iraq

*Corresponding

author:

munamm_math@csw.uobaghdad.edu.iq

*ORCID ID: 0000-0001-8620-4976

optimal solution of the latest problem as an approximate one with a controllable error for the original solution is obtained (6). Nazemi and Farahi in 2011 considered a numerical method for nonlinear Fredholm integral equations of the second kind with the continuous kernel by converting the integral equation problem into an optimization problem (7). Skandari et al. in 2011 proposed a new approach for a class of optimal control problems to solve Volterra integral equations which is based on linear combination property of intervals (8). Erfanian and Mostahsan in 2011 considered two stages of approximation to find an approximate solution for a class of nonlinear Volterra integral equations by: first convert the integral equation to a moment problem, and then modify the new problem to two classes of optimization problems (non-constraint optimization and optimal control problems) (9). Effati and Skandari in 2012 presented a new approach for linear Volterra integral equations based on optimal control theory and some optimal control problems corresponding Volterra integral equation are introduced which solved by discretization methods and linear programming approaches (10).

Many authors have paid attention for studying the numerical solution of LVFIE. Ahmed in 2011 investigated the numerical solution for a mixed linear Volterra-Fredholm integral equation by

approximating the unknown function using a tensor product (Algebraic or Chebyshev)-surface and substituting it in the Volterra-Fredholm integral equations, second apply least-square technique for minimizing the error terms on the given domain, and then obtain a system of linear algebraic equations which solved using control points (11). Chen and Jiang in 2012 used Taylor expansion method to solve a mixed linear Volterra-Fredholm integral equation of the 2nd kind (12). Ghanim in 2014 introduced numerical algorithms to find numerical solution for linear Volterra – Fredholm integral equation of the 2nd kind using three different kinds of Lagrange polynomials (13). Mustafa and Harbi in 2014 used non-polynomial spline function to find numerical solution for Volterra integral equation of the 2nd kind (14). Nemati et al. in 2015 proposed numerical solution of linear mixed Volterra-Fredholm integral equations in one space variable, the proposed numerical algorithm combines the trapezoidal rule, for the integration in time, with piecewise polynomial approximation, for the space discretization (15). Chniti and Alhazmi in 2015 presented smoothing transformation, Legendre and Chebyshev collocation method to solve numerically the Volterra-Fredholm integral equations with logarithmic kernel (16). Khan et al. in 2017 provided a numerical technique for obtaining approximate solution of mixed Volterra-Fredholm integral equations of 2nd kind based on the Bernstein’s approximation (17). Hasan et al. in 2017 proposed a fixed point method to solve a system of LVFIE of the 2nd kind using fixed point method (18).

In this paper, LVFIE of the 2nd kind of the following form are considered:

$$g(x) = f(x) + \int_a^x w_1(x,t)g(t)dt + \int_a^b w_2(x,t)g(t)dt, \quad x \in [a, b] \quad \dots (1)$$

where $g(x)$ is the unknown function to be determined, $f(x), w_1(x, t)$ and $w_2(x, t)$ are continuous known functions. Therefore an approximate solution depending on a polynomial of degree $(m - 1)$ is proposed, which has the form:

$$g(x) \approx g_m(x) = \sum_{i=1}^m c_i x^{i-1} \quad \dots (2)$$

where c_1, c_2, \dots, c_m are arbitrary constants to be determined. Next using numerical integration includes (TR, SR, BR, and RI) to find an approximate solution for the integral parts in Eq.1, after that using LPP to find an approximate solution to this problem. This paper is well ordered as follows: section (2) includes basic definitions, section (3) contains transforms Eq.1 to the LPP, in section (4), the algorithm for solving LVFIE is proposed, section (5) includes test examples.

Finally, section (6) gives conclusions and recommendations.

Basic Definitions:

Definition 1: An integral equation can be recognized as equations in which the indefinite function $g(x)$ to be specific appear under the integral sign. A standard formula of an integral equation in $g(x)$ is:

$$c(x)g(x) = f(x) + \lambda \int_{a(x)}^{b(x)} w(x,t)g(t) dt, \quad x \in [a(x), b(x)] \quad \dots (3)$$

where $c(x), f(x)$ and $w(x, t)$ are given continuous functions, $w(x, t)$ is known as the kernel of the integral Eq. 3, and $a(x)$ and $b(x)$ are the limits of integration, λ is a constant parameter (19). Also, Eq. 3 is called ‘Volterra integral equation’ (VIE), if $b(x) = x$ and $a(x)=a$, and it is called ‘Fredholm integral equation’ (FIE), if $b(x) = b$ and $a(x) = a$, where a and b are constants.

Note that, for Eq. 1 when $w_1(x, t)$ is equal to zero Eq. 1 becomes FIE, furthermore when $w_2(x, t)$ is equal to zero Eq. 1 becomes VIE. And it is called VFIE if the both integral appear at one time. In this paper Eq. 1 is taken as a general form to improve that the proposed method is effective for the three kinds of equations.

Definition 2: The general form for LPP can be expressed mathematically as follows (20):

$$\left. \begin{array}{l} \text{minimize or maximize } z = \sum_{j=1}^n c_j x_j \\ \text{Subject to} \\ \sum_{j=1}^n a_{ij} x_j (\leq, =, \geq) b_i, \quad i = 1, 2, \dots, m \\ x_j \geq 0, \quad j = 1, 2, \dots, n \end{array} \right\} \quad \dots (4)$$

where c_j ($j = 1, \dots, n$) are real number called cost coefficients, b_i ($i = 1, \dots, m$) are real number called stipulations, a_{ij} ($i = 1, 2, \dots, m; j = 1, 2, \dots, n$) are constants called structural coefficients and x_j ’s are called decision variables for ($j = 1, 2, \dots, n$).

Transform the LVFIE’s of the 2nd Kind to LPP:

The solution of LVFIE of the 2nd kind can be founded by substituting $g_m(x)$ in the Eq. 2 instead of the unknown function in Eq. 1 yields:

$$\sum_{i=1}^m c_i x^{i-1} = f(x) + \int_a^x w_1(x,t) \sum_{i=1}^m c_i t^{i-1} dt + \int_a^b w_2(x,t) \sum_{i=1}^m c_i t^{i-1} dt + \epsilon(x)$$

where $\epsilon(x)$ is the approximate error, then we get:

$$\epsilon(x) = f(x) - \sum_{i=1}^m c_i x^{i-1} + \int_a^x w_1(x,t) \sum_{i=1}^m c_i t^{i-1} dt + \int_a^b w_2(x,t) \sum_{i=1}^m c_i t^{i-1} dt \quad \dots (5)$$

Simplification of Eq. 5, to obtain:

$$\epsilon(x) = f(x) - \sum_{i=1}^m c_i \left[x^{i-1} - \int_a^x w_1(x,t)t^{i-1} dt - \int_a^b w_2(x,t)t^{i-1} dt \right] \dots \quad (6)$$

Suppose

$$\psi_i(x) = x^{i-1} - \int_a^x w_1(x,t)t^{i-1} dt - \int_a^b w_2(x,t)t^{i-1} dt, \quad i = 1, 2, \dots, m \dots \quad (7)$$

Now, substituting Eq. 7 in Eq. 6 yields:

$$\epsilon(x) = f(x) - \sum_{i=1}^m c_i \psi_i(x) \dots \quad (8)$$

and dividing the interval $[a, b]$ into n subintervals to get the points $x_j = a + (j - 1) * h, j = 1, 2, \dots, n + 1$ where $h = \frac{b-a}{n}, x_1 = a,$ and $x_{n+1} = b$. Hence, after substituting $x = x_j,$ for $j = 1, 2, \dots, n + 1$ in Eq. 7, the following formula is obtained:

$$\psi_i(x_j) = \begin{cases} x_j^{i-1} - \int_a^{x_j} w_1(x,t)t^{i-1} dt & , \quad \text{if } i = 1 \\ x_j^{i-1} - \int_a^{x_j} w_1(x,t)t^{i-1} dt - \int_a^b w_2(x,t)t^{i-1} dt & , \quad \text{if } 2 \leq i \leq m \end{cases}, \quad j = 1, 2, \dots, n + 1 \dots \quad (9)$$

Therefore, substituting $x = x_j$ for $j = 1, 2, \dots, n + 1,$ Eq. 8 becomes:

$$\epsilon(x_j) = f(x_j) - \sum_{i=1}^m c_i \psi_i(x_j), \quad j = 1, 2, \dots, n + 1 \dots \quad (10)$$

Now, define γ as:

$$\gamma = \sum_{j=1}^{n+1} \epsilon(x_j) \dots \quad (11)$$

Therefore, Eq. 1 can be written in the following LPP form:

$$\left. \begin{aligned} \min \gamma &= \sum_{j=1}^{n+1} \epsilon(x_j) \\ \text{s.t} & \\ & \left. \begin{aligned} \sum_{i=1}^m c_i \psi_i(x_j) - \gamma &\leq f(x_j) \\ - \sum_{i=1}^m c_i \psi_i(x_j) - \gamma &\leq -f(x_j) \end{aligned} \right\} j = 1, \dots, n + 1 \end{aligned} \right\} \dots \quad (12)$$

where c_i 's are unrestricted in sign, $\psi_i(x_j)$'s can be estimated numerically as shown in the following subsection. Hence, the values of c_i 's are used to get the approximate solution $g_m(x)$ in Eq. 2, which is the approximate solution of LVFIE of the 2nd kind [i.e. Eq. 1].

Calculate $\psi_i(x_j)$ Using Numerical Integration Formulas:

In this subsection, the numerical computation of the integral part in $\psi_i(x_j)$ is shown using quadrature methods including (TR, SR, BR, and RI). For this purpose, the following notation is used: $\int_{a_1}^{b_1} w(x_j, t)t^{i-1} dt$ for denotation of the integral part in $\psi_i(x_j), j = 1, 2, \dots, n + 1.$

Suppose $a_1 = t_1 < t_2 < \dots < t_n = b_1, t_k = t_1 + (k - 1)h, k = 1, 2, \dots, n + 1,$ and $h = \frac{b_1 - a_1}{n}$

1. Using TR:

$$\int_{a_1}^{b_1} w(x_j, t)t^{i-1} dt = \frac{h}{2} [w(x_j, a_1)a_1^{i-1} + 2 \sum_{k=2}^n w(x_j, t_k)t_k^{i-1} + w(x_j, b_1)b_1^{i-1}] \dots \quad (13)$$

2. Using SR:

$$\int_{a_1}^{b_1} w(x_j, t)t^{i-1} dt = \frac{h}{3} [w(x_j, a_1)a_1^{i-1} + 4w(x_j, t_2)t_2^{i-1} + 2w(x_j, t_3)t_3^{i-1} + 4w(x_j, t_4)t_4^{i-1} + \dots + 4w(x_j, t_n)t_n^{i-1} + w(x_j, b_1)b_1^{i-1}] \dots \quad (14)$$

3. Using BR:

$$\int_{a_1}^{b_1} w(x_j, t)t^{i-1} dt = \frac{2h}{45} [7w(x_j, a_1)a_1^{i-1} + 32w(x_j, t_2)t_2^{i-1} + 12w(x_j, t_3)t_3^{i-1} + 32w(x_j, t_4)t_4^{i-1} + 14w(x_j, t_5)t_5^{i-1} + \dots + 32w(x_j, t_n)t_n^{i-1} + 7w(x_j, b_1)b_1^{i-1}] \dots \quad (15)$$

4. Using RI:

$$\int_{a_1}^{b_1} w(x_j, t)t^{i-1} dt = \frac{h}{2} \left[w(x_j, a_1)a_1^{i-1} + 2 \sum_{k=2}^{2^j-1} w(a_1 + (2k - 1) * h_j, t_k)t_k^{i-1} + w(x_j, b_1)b_1^{i-1} \right] \dots \quad (16)$$

The Algorithm for Solving LVFIE of the 2nd kind:

The following steps are used to find an approximate solution of LVFIE of the 2nd kind:

Step 1. Select two positive integers m and $n,$ (where m represent the number of terms of series in Eq. 2, n represent the number of subinterval for the closed interval $[a,b]$).

Step 2. Calculate $f(x_j), j = 1:n + 1.$

Step 3. Using numerical integration to calculate $\psi_i(x_j)$ in Eq. 8 for each $i = 1, 2, \dots, m; j = 1, 2, \dots, n + 1,$ which are (TR, SR, BR, and RI).

i. TR, by using Eq. 13.

ii. SR, by using Eq. 14.

iii. BR, by using Eq. 15.

iv. RI, by using Eq. 16.

Step 4. Compute γ from Eq. 11.

Step 5. Construct the LP model Eq. 12.

Step 6. Using generalized simplex method to find the optimal approximation of $c_i, i = 1, 2, \dots, m.$

Step 7. Determine the approximate solution of Eq. 1 by substituting the values of $c_i, i = 1, 2, \dots, m$ in Eq. 2.

Note that in this paper MATLAB R2018a is used for implementation of the algorithm.

Numerical Test Examples:

In this section, some of the numerical test examples are given to illustrate the proposed method for solving the LVFIE of the 2nd kind. In all the test examples $g(x)$ is chosen in such a way that we know the exact solution. The exact solution is used only to show that the numerical solution obtained with our method is true. Then, in these test

examples the absolute error are calculated in all points.

Test Example 1: Consider the LVFIE of 2nd kind (17):

$$g(x) = x - 2e^x + e^{-x} + 1 + \int_0^x te^x g(t) dt + \int_0^1 e^{x+t} g(t) dt$$

for which the exact solution is $g(x) = e^{-x}$. Table 1 shows the absolute error obtained by using (TR, SR, BR, and RI) for $h = 0.1, n = 10, m = 10$, where m is the degree of approximate polynomial $g(x)$ appear in Eq. 2 and $\|err\|_\infty$ is the maximum absolute error, for $x \in [0,1]$.

Table 1. The absolute error of test example 1

x	Error using TR	Error using SR	Error using BR	Error using RI
0	0.000102493241609158	2.49109375349832e-08	2.01430538915304e-10	2.27079022252497e-10
0.1	0.000113451577860246	2.77139680093796e-08	1.86916038202867e-10	2.15425566274519e-10
0.2	0.000124537294404647	3.11356900262183e-08	1.98265071027492e-10	2.30316765659211e-10
0.3	0.000133715127830891	3.68140724571475e-08	1.37722810933383e-09	1.34068800505815e-09
0.4	0.000138651792887523	3.9765073456266e-08	1.18836496199037e-10	1.60952806638193e-10
0.5	0.00013651617237409	4.35975450185921e-08	1.32601707392155e-10	1.81023751544274e-10
0.6	0.000123448433075213	4.56720178343417e-08	7.85884357590305e-10	7.32290228455668e-10
0.7	9.37002095329209e-05	3.9269196672187e-08	1.74255054830041e-11	6.94454493910257e-11
0.8	3.80499682491875e-05	1.93316483931838e-08	7.09133862741851e-11	1.00750630060986e-10
0.9	5.94811463363598e-05	2.88283432681169e-08	7.03646030331129e-11	2.69834155020021e-11
1	0.000228347336972523	1.32361363092137e-07	6.38649244599776e-10	4.09659972611109e-10
$\ err\ _\infty$	0.000228347336972523	1.32361363092137e-07	1.37722810933383e-09	1.34068800505815e-09

Table 2 shows the maximum absolute error of test example 1 by using RI with 10 columns for $n = 5, 10, \dots, 30$ and $m = 5, 10, \dots, 20$ and

comparing with the minimum error in (17) obtained using Bernstein polynomials.

Table 2. Maximum absolute error by using RI with 10 columns

	m=5	m=10	m=15	m=20	Minimum error in (17)
n=5	1.80466052200901e-05	1.73236324783055e-05	0.000139531997039199	0.00048852860377735	
n=10	1.29375330428783e-05	1.34068800505815e-09	2.30496499664667e-09	2.08972145032682e-07	
n=15	1.2817840279622e-05	1.83188264557543e-09	5.94055804548077e-10	3.00607217784687e-07	7.071067812*
n=20	1.17171123399373e-05	1.86609017127637e-09	4.58927396085329e-09	1.19387566610563e-07	10⁻¹¹
n=25	1.12564382384051e-05	2.68936872771519e-09	3.24708193666368e-09	1.97195315632115e-09	
n=30	1.11584049335978e-05	1.78146986051786e-09	5.00741970110852e-09	1.49425165574257e-08	

Test Example 2: Consider the FIE of the 2nd kind (21): $g(x) = \sin(x) - x + \int_0^{\pi/2} xt g(t) dt$ for which the exact solution is $g(x) = \sin(x)$.

Table 3 shows the absolute error obtained by using (TR, SR, BR, and RI) for $h = 0.1, n = 10, m = 10$, and $x \in [0, \frac{\pi}{2}]$.

Table 3. The absolute error of test example 2

x	Error using TR	Error using SR	Error using BR	Error using RI
0	0	0	0	0
$\frac{\pi}{20}$	0.000275096664244356	3.46922592708854e-07	3.93723234570764e-09	5.28284549439206e-09
$\frac{2\pi}{20}$	0.000550193328488546	6.84035879827682e-07	1.93483751242951e-09	7.5638911800624e-10
$\frac{3\pi}{20}$	0.000825955324154082	1.0238570866683e-06	5.09898773204398e-09	1.06214725903442e-09
$\frac{4\pi}{20}$	0.00110038665697776	1.3680717573239e-06	3.86967580201514e-09	1.51277890214629e-09
$\frac{5\pi}{20}$	0.00137482378285103	1.7118563248042e-06	3.070467546884e-09	3.65760144394045e-09
$\frac{6\pi}{20}$	0.00165057998546692	2.05210763648545e-06	5.80451475773458e-09	2.26916840873059e-09
$\frac{7\pi}{20}$	0.00192639040596387	2.39212967390046e-06	8.76783445846741e-09	6.5146210648237e-10
$\frac{8\pi}{20}$	0.00220077331395585	2.73614351353757e-06	7.73935204811949e-09	3.02555758224798e-09
$\frac{9\pi}{20}$	0.00247586997820104	3.07816145295181e-06	8.70677108189e-09	3.40375128082826e-09
$\frac{\pi}{2}$	0.00275024709314409	3.41770074263614e-06	1.21528355156997e-08	1.30330013448088e-09
$\ err\ _\infty$	0.00275024709314409	3.41770074263614e-06	1.21528355156997e-08	5.28284549439206e-09

Table 4 shows the maximum absolute error of test example 2 by using RI with 10 columns for $n = 5, 10, \dots, 30$ and $m = 5, 10, \dots, 20$ and

comparing with the minimum error in (21) using A wavelet based method.

Table 4. Maximum absolute error of test example 2

	m=5	m=10	m=15	m=20	Minimum error in (21)
n=5	0.000250532304200357	0.00204524171715315	0.00700937506399091	0.00299344393818657	4.87051*10⁻⁷
n=10	0.000150493052866185	5.28284549439206e-09	2.95809066130914e-09	1.36430372665863e-07	
n=15	0.000126432937772925	1.25692456443005e-10	3.26260370886455e-07	2.07978079203031e-12	
n=20	0.000122723915813472	9.42873719211867e-07	5.11214459653075e-09	7.28038746622417e-07	
n=25	0.000134526659284484	1.53481720666093e-08	3.21489566856847e-07	5.01681764353279e-07	
n=30	0.000130829048964776	7.12419012671717e-11	3.28022662254845e-07	6.90780765921772e-13	

Test Example 3: Consider the VIE of the 2nd kind. (22)

error obtained by using (TR, SR, BR, and RI) for $h = 0.1, n = 10, m = 10$, and $x \in [0, 1]$.

$g(x) = x^5 - \frac{x^8}{7} + \int_0^x xtg(t)dt$ for which the exact solution is $g(x) = x^5$. Table 5 shows the absolute

Table 5: The absolute error of test example 3

x	Error using TR	Error using SR	Error using BR	Error using RI
0	0	0	0	0
0.1	4.07240313671209e-08	8.24038275934354e-08	4.7171878984503e-10	3.3881317890172e-21
0.2	5.14835299653517e-08	1.63159514236203e-07	9.34002835088679e-10	1.0842021724855e-19
0.3	1.0280117092613e-07	2.34026355174985e-07	1.33967841290869e-09	8.67361737988404e-19
0.4	8.117302906889e-07	2.73578518049725e-07	1.56609384314998e-09	3.46944695195361e-18
0.5	4.94217156043336e-06	2.40612479093794e-07	1.37738051173009e-09	1.38777878078145e-17
0.6	2.14923150374663e-05	6.75544593181643e-08	3.86713883138157e-10	2.77555756156289e-17
0.7	7.45893556647337e-05	3.46132139256161e-07	1.98142546814495e-09	5.55111512312578e-17
0.8	0.000220789721384063	1.1405392973951e-06	6.52899079245373e-09	1.11022302462516e-16
0.9	0.000579980666897018	2.50190694273122e-06	1.43221082771205e-08	3.33066907387547e-16
1	0.00138699622275107	4.66921551289801e-06	2.6728815827326e-08	4.44089209850063e-16
$\ err\ _\infty$	0.00138699622275107	4.66921551289801e-06	2.6728815827326e-08	4.44089209850063e-16

Table 6 shows the maximum absolute error of test example 3 by using RI with 10 columns with $n = 5, 10, \dots, 30$ and $m = 5, 10, \dots, 20$ and

comparing with the minimum error in (22) using analytical techniques for a numerical solution.

Table 6. Maximum absolute error of test example 3

	m=5	m=10	m=15	m=20	Minimum error in (22)
n=5	0.00344432832521402	3.33066907387547e-16	0.00859839736540469	0.0236481084743722	4.63*10⁻¹⁰
n=10	0.00210910706319622	4.44089209850063e-16	1.01918473660589e-13	2.28582959427293e-06	
n=15	0.00219352883926544	8.21565038222616e-15	5.77315972805081e-15	4.37383462781327e-12	
n=20	0.00203247879180546	1.38222766565832e-13	7.21644966006352e-15	1.09923181668137e-12	
n=25	0.00206776701090695	7.105427357601e-14	7.7715611723761e-15	5.313527395856e-13	
n=30	0.00199598890700549	3.84636766881385e-13	5.46229728115577e-14	4.3310910413652e-12	

Conclusions:

This paper presents a method of finding the solution of LVFIE of the 2nd kind using the LPP. The polynomial of degree $m - 1$ is used to convert the LVFIE of the 2nd kind into LPP. TR, SR, BR and RI are proposed for computing the integral part and the comparison between those methods is made. The accuracy of the method has been shown by applying different test examples and comparing the results with the exact solution. The results for the LPP are improved using RI instead of the other methods to evaluate the integrals within linear

programming method. Also, it is supposed that the best result can be obtained by increasing both the number of basic functions (m) and the number of constraints (n) with keeping ($n > m$). For future work, we suggest using this method for solving ‘Volterra- Fredholm integro-differential equations’, by using suitable approximation for derivatives part.

Conflicts of Interest: None.

References:

- Kanwal RP. Linear Integral Equations Theory and Techniques. New York: Academic Press, INC.; 1971. 296 p.
- AL-Nasir RH. Numerical Solution of Volterra Integral Equations of the Second Kind. M. Sc. Thesis. Baghdad: University of Technology; 1999.
- Saed NS. Numerical Solution of Fredholm Integral Equations of the First Kind. M. Sc. Thesis. Baghdad: University of Technology; 1999.
- Kalwi PM. Numerical Methods for Solving Fredholm Integral Equations of The Second Kind. M. Sc. Thesis. Baghdad: University of Technology; 1999.
- Shua'a AH. Using Linear Programming to find solution Volterra Integral Equation. J of Col of B ED. 2005;7(43):12-31.
- Kamyad AV, Mehrabinezhad M, Sabri-Nadjafi J. A Numerical Approach for Solving Linear and Nonlinear Volterra Integral Equations with Controlled Error. IAENG IJAM. 2010; 40(2): 69-74.
- Nazemi AR, Farahi MH. A Numerical Scheme for Fredholm Integral Equations. JARAM. 2011; 3(1): 47-62.
- Skandari MH, Erfanian HR, Kamyad AV. A New Approach for a Class of Optimal Control Problems of Volterra Integral Equations. ICA. 2011; 2: 121-125.
- Erfanian HR, Mostahsan T. Approximate Solution of a Class of Nonlinear Volterra Integral Equations. J Math Computer Sci. 2011; 3(3): 278-286.
- Effati S, Skandari MH. Optimal Control Approach for Solving Linear Volterra Integral Equations. IJISA. 2012; 4: 40-46.
- Ahmed ShSh. Numerical Solution for Volterra-Fredholm Integral Equation of the Second Kind by Using Least Squares Technique. IJS. 2011; 52(4): 504-512.
- Chen Z, Jiang W. An Approximate Solution for a Mixed Linear Volterra-Fredholm Integral Equation. App Math Lett. 2012; 25: 1131-1134.
- Ghanim IN. Modified Numerical Algorithms for Solving Linear Volterra-Fredholm Integral Equations of the Second Kind". M. Sc. Thesis. Baghdad: University of Baghdad; 2014.
- Mustafa MM, Harbi SH. Solution of Second Kind Volterra Integral Equations Using Non-Polynomial Spline Functions. Baghdad Sci J. 2014; 11(2): 207-210.
- Nemati S, Lima P, Ordokhani Y. Numerical Method for the Mixed Volterra-Fredholm Integral Equations Using Hybrid Legendre Functions. In: Brandts J, Korotov S, Křížek M, Segeth K, Šístek J, Vejchodský T, editors. Conference Applications of Mathematics, in honor of the birthday anniversaries of Ivo Babuška (90), Milan Práger (85), and Emil Vitásek (85). Prague: Institute of Mathematics AS CR; 2015: 184-193.
- Chniti Ch, Alhazmi ShA. On the Numerical Solution of Volterra-Fredholm Integral Equations with Logarithmic Kernel Using Smoothing Transformation. IJAMR. 2015; 4 (1): 183-192.
- Khan F, Mustafa G, Omar M, Komal H. Numerical Approach Based on Bernstein Polynomials for Solving Mixed Volterra-Fredholm Integral Equations. AIP Advances. 2017; 7(12): ID 125123.
- Hasan TI, Salleh Sh, Sulaiman NA. Fixed Point Method and its Improvement for the System of Volterra-Fredholm Integral Equations of the Second Kind. Matematika. 2017; 33(2): 191-206.
- Wazwaz AM. Linear and Nonlinear Integral Equations Methods and Applications. Beijing: Higher Education Press and Berlin Heidelberg: Springer-Verlag; 2011. 639 p.
- Gupta PK, Hira DS. Operation Research. 7th Revised Edition. New Delhi: S. CHAND & COMPANY PVT. LTD; 2014. 1498 p.
- Lin En-Bing, Al-Jarrah Y. A Wavelet Based Method for the Solution of Fredholm Integral Equations. AJCM. 2012; 2(2): 114-117.
- Berenguer MI, Gámez D, Garralda-Guillem AI, Ruiz Galón M, Serrano Pérez MC. Analytical Techniques for a Numerical Solution of the Linear Volterra Integral Equation of the Second Kind. Abs Appl Anal. 2009; 2009: Article ID 149367.

حل معادلة فولتيرا- فريدهولم التكاملية الخطية من النوع الثاني باستخدام طريقة البرمجة الخطية

منى منصور مصطفى

نجوى رحيم مصطفى

ایمان عبد الهادي ضاري

قسم الرياضيات، كلية العلوم للبنات، جامعة بغداد، بغداد، العراق.

الخلاصة:

في هذا البحث تم عرض تقنية جديدة لإيجاد حل لثلاثة أنواع من المعادلات التكاملية الخطية من النوع الثاني المتضمنة: معادلة فولتيرا- فريدهولم التكاملية (LVFIE) (الحالة العامة)، معادلة فولتيرا التكاملية (LVIE) و معادلة فريدهولم التكاملية (LFIE) (كحالتين خاصتين). التقنية الجديدة تعتمد على تقريب الحل الى متعددة حدود من الدرجة (m-1) وبعد ذلك تحويل المسألة الى مسألة برمجة خطية (LPP) والتي سوف تحل لإيجاد الحل التقريبي لمعادلة فولتيرا- فريدهولم التكاملية الخطية من النوع الثاني (LVFIE). علاوة على ذلك تم استخدام الطرق التريبيعية التي تضم: قاعدة شبه المنحرف (TR)، قاعدة سمبسون 1/3 (SR)، قاعدة بول (BR) وصيغة رومبرك للتكامل (RI) لتقريب التكامل الموجود في LVFIE، كما تم عمل مقارنات بين هذه الطرق. واخيرا لزيادة التوضيح تم اعطاء الخوارزمية المتبعة في الحل وتم تطبيقها على امثلة اختبارية لتوضيح فعالية التقنية الجديدة.

الكلمات المفتاحية: معادلة فولتيرا- فريدهولم التكاملية الخطية من النوع الثاني، مسألة البرمجة الخطية، الطرق التريبيعية.