



ISSN: 0067-2904

A Generalized Subclass of Starlike Functions Involving Jackson's (p, q) -Derivative

Abdeljabbar Talal Yousef*, Zabidin Salleh

School of Informatics and Applied Mathematics, University Malaysia Terengganu, 21030 Kuala Nerus, Terengganu, Malaysia.

Received: 9/6/ 2019

Accepted: 21/ 9/2019

Abstract

In this paper, we generalize many earlier differential operators which were studied by other researchers using our differential operator. We also obtain a new subclass of starlike functions to utilize some interesting properties.

Keywords: Differential operator, starlike functions, coefficient inequality, inclusion properties, convexity.

1. Introduction

Let A represents the class of all analytic functions φ defined in the open unit disk $\Theta = \{z \in \mathbb{C} : |z| < 1\}$, and normalized by the conditions $\varphi(0) = 0$ and $\varphi'(0) = 1$. Therefore, each $\varphi \in A$ has a Taylor-Maclaurin series extension of the form:

$$\varphi(z) = z + \sum_{h=2}^{\infty} a_h z^h, (z \in \Theta) \quad (1.1)$$

Furthermore, let S represents the class of all functions $\varphi \in A$ which are univalent in Θ . The quantum calculus (henceforth q -calculus) is considered as a crucial tool that is used to explore the subclasses of analytic functions. q -calculus operators were used by Kanas and Raducanu to investigate some significant classes of functions which are analytic in Θ [1]. The importance of the fractional calculus applications is obvious in many topics of mathematics, such as in the fields of q -transform analysis, ordinary fractional calculus, and operator theory. Recently, researchers paid more attention to the area of q -calculus and several new operators have been proposed. The application of q -calculus was first founded by Jackson who developed the q -integral and q -derivative in a systematic way [2]. After that, through several studies on quantum groups, the geometrical interpretation of q -analysis was identified. Unlike the typical calculus, this calculus has no limits notion. A good detailed work on the calculus and its applications in operator theory is found in a previous report [3], while more information were provided in other articles [4, 5].

The main structure of (p, q) -calculus was established on only one parameter, but since then it was generalized to the post-quantum calculus (represented by (p, q) -calculus). In this section, we assume that we can obtain calculus by substituting $p = 1$ in calculus.

To be fulfilled, some brief notations and definitions of (p, q) -calculus are provided below: For Jackson's derivative where $0 < p < q \leq 1$ and $\varphi \in A$, the following is provided [2]:

*Email: abduljabaryousef@gmail.com

$$D_{p,q}\varphi(z) = \begin{cases} \frac{\varphi(pz) - \varphi(qz)}{(p-q)z} & \text{for } z \neq 0. \\ \varphi'(0) & \text{for } z = 0. \end{cases} \quad (1.2)$$

From (1.2), we have

$$D_{p,q}\varphi(z) = 1 + \sum_{h=2}^{\infty} [h]_{p,q} a_h z^{h-1} \quad (1.3)$$

Where

$$[h]_{p,q} = p^{h-1} + p^{h-2}q + p^{h-3}q^2 + \dots + pq^{h-2} + q^{h-1} = \frac{p^h - q^h}{p - q} \quad (1.4)$$

is named (p, q) -bracket. It's notable that when $p = 1$, the bracket is an obvious generalization of the q -number, that is

$$[h]_{1,q} = \frac{1 - q^h}{1 - q} = [h]_q, q \neq 1$$

For $p = 1$, one can notice that the Jackson's (p, q) -derivative will be reduced to the q -derivative, as previously described [2]. It was clearly proved that for a function $\gamma(z) = z^h$, the

$D_{p,q}\gamma(z) = D_{p,q}z^h = \frac{p^h - q^h}{p - q} z^{h-1} = [h]_{p,q} z^{h-1}$ is obtained. For $\varphi \in A$, the

Sălăgean (p, q) -differential operator is defined as follows [6]:

$$\begin{aligned} \Gamma_{p,q}^0 \varphi(z) &= \varphi(z), \\ \Gamma_{p,q}^1 \varphi(z) &= z D_{p,q} \varphi(z), \\ &\dots \\ \Gamma_{p,q}^k \varphi(z) &= \Gamma_{p,q}^1 (\Gamma_{p,q}^{k-1} \varphi(z)), \\ &= z + \sum_{h=2}^{\infty} [h]_{p,q}^k a_h z^h, \quad (k \in \mathbb{N}_0 = \mathbb{N} \cup \{0\}, z \in \Theta) \end{aligned} \quad (1.5)$$

It's observable when $p = 1$ and $\lim_{q \rightarrow 1^-}$, the well-known Sălăgean operator is obtained [6]:

$$\Gamma^k \varphi(z) = z + \sum_{h=2}^{\infty} h^k a_h z^h, \quad (z \in \Theta) \quad (1.6)$$

Now let

$$\begin{aligned}
\Lambda_{\beta,\delta,\lambda,p,q}^{0,k} \varphi(z) &= \Gamma_{p,q}^k \varphi(z), \\
\Lambda_{\beta,\delta,\lambda,p,q}^{1,k} \varphi(z) &= (1 - \beta(\delta - \lambda)) \Gamma_{p,q}^k \varphi(z) + \beta(\delta - \lambda) z \left(\Gamma_{p,q}^k \varphi(z) \right)' \\
&= z + \sum_{h=2}^{\infty} [h]_{p,q}^k [1 + \beta(\delta - \lambda)(h-1)] a_h z^h. \\
\Lambda_{\beta,\delta,\lambda,p,q}^{2,k} \varphi(z) &= (1 - \beta(\delta - \lambda)) \Lambda_{\beta,\delta,\lambda,p,q}^{1,k} \varphi(z) + \beta(\delta - \lambda) z \left(\Lambda_{\beta,\delta,\lambda,p,q}^{1,k} \varphi(z) \right)' \\
&= z + \sum_{h=2}^{\infty} [h]_{p,q}^k [1 + \beta(\delta - \lambda)(h-1)]^2 a_h z^h.
\end{aligned} \tag{1.7}$$

In general, we have

$$\begin{aligned}
\Lambda_{\beta,\delta,\lambda,p,q}^{\zeta,k} \varphi(z) &= (1 - \beta(\delta - \lambda)) \Lambda_{\beta,\delta,\lambda,p,q}^{\zeta-1,k} \varphi(z) + \beta(\delta - \lambda) z \left(\Lambda_{\beta,\delta,\lambda,p,q}^{\zeta-1,k} \varphi(z) \right)' \\
&= z + \sum_{h=2}^{\infty} [h]_{p,q}^k [1 + \beta(\delta - \lambda)(h-1)]^{\zeta} a_h z^h
\end{aligned} \tag{1.8}$$

Where $\beta \geq 0, \lambda \geq 0, \delta \geq 0$ and $\zeta \in \mathbb{N}_0$.

It is observable that we have $\Lambda_{\beta,\delta,\lambda,p,q}^{0,0} \varphi(z) = \varphi(z)$, and $\Lambda_{\beta,\delta,\lambda,p,q}^{1,0} \varphi(z) = z \varphi'(z)$. It is noticeable that when $p=1$, the differential operator $\Lambda_{\beta,q}^{\zeta,k} \varphi(z)$ that was defined and studied by Frasin and Murugusundaramoorthy is obtained [7]. Also, it is noticeable that when $p=1$ and $\lim_{q \rightarrow 1}$, the following differential operator is obtained:

$$\Lambda_{\beta}^{\zeta,k} \varphi(z) = z + \sum_{h=2}^{\infty} h^k [1 + \beta(\delta - \lambda)(h-1)]^{\zeta} a_h z^h$$

It is noticeable that when $\delta=1$ and $\lambda=0$, we find the differential operator $\Lambda_{\beta,p,q}^{\zeta,k} \varphi(z)$ that was defined and studied by Feras Yousef [8]. Furthermore, when $k=0$ we find the differential operator $\Lambda_{\beta,\delta,\lambda}^{\zeta}$ that was defined and studied by Ibrahim and Darus [9, 10], and when $\delta=1$, $\lambda=0$ and $k=0$ we identify the differential operator Λ_{β}^{ζ} defined and studied by Al-Oboudi [10], while if $\zeta=0$, we identify Sălăgean differential operator Λ^{ζ} [6].

By using the differential operator $\Lambda_{\beta,\delta,\lambda,p,q}^{\zeta,k} \varphi(z)$, we say that a function $\varphi(z)$ belonging to A is in the class $\mathcal{Q}_{p,q}^{\zeta,k}(\beta, \delta, \lambda, \mu, b)$ if and only if

$$\begin{aligned}
&\left| \frac{(1 - \beta(\delta - \lambda)) z \left(\Lambda_{\beta,\delta,\lambda,p,q}^{\zeta,k+1} \varphi(z) \right)' + \beta(\delta - \lambda) z \left(\Lambda_{\beta,\delta,\lambda,p,q}^{\zeta,k+1} \varphi(z) \right)}{(1 - \beta(\delta - \lambda)) \Lambda_{\beta,\delta,\lambda,p,q}^{\zeta,k} \varphi(z) + \beta(\delta - \lambda) \Lambda_{\beta,\delta,\lambda,p,q}^{\zeta,k} \varphi(z)} - 1 \right| \\
&< \mu \left| \frac{(1 - \beta(\delta - \lambda)) z \left(\Lambda_{\beta,\delta,\lambda,p,q}^{\zeta,k+1} \varphi(z) \right)' + \beta(\delta - \lambda) z \left(\Lambda_{\beta,\delta,\lambda,p,q}^{\zeta,k+1} \varphi(z) \right)}{(1 - \beta(\delta - \lambda)) \Lambda_{\beta,\delta,\lambda,p,q}^{\zeta,k} \varphi(z) + \beta(\delta - \lambda) \Lambda_{\beta,\delta,\lambda,p,q}^{\zeta,k} \varphi(z)} - b \right|, \quad (k, \zeta \in \mathbb{N}_0)
\end{aligned} \tag{1.9}$$

for some $\mu(0 \leq \mu < 1)$, $\beta, \delta, \lambda \geq 0$, and $0 \leq b < 1$ for all $z \in \Theta$.

Let T denotes the subclass of A consisting of functions of the form

$$\varphi(z) = z - \sum_{h=2}^{\infty} a_h z^h \quad (a_h \geq 0, z \in \Theta) \tag{1.10}$$

Further, we define the class $P_{p,q}^{\zeta,k}(\beta, \delta, \lambda, \mu, b)$ by

$$P_{p,q}^{\zeta,k}(\beta, \delta, \lambda, \mu, b) = Q_{p,q}^{\zeta,k}(\beta, \delta, \lambda, \mu, b) \cap T$$

The main target of this paper is to provide a systematic investigation of some important features and characteristics of the class $P_{p,q}^{\zeta,k}(\beta, \delta, \lambda, \mu, b)$. Some interesting corollaries and natural consequences of the main findings are also considered. Some important techniques used earlier by many researchers were applied in this work (see Al-Hawary et al. [11, 12], Aouf and Srivastava [13], and Frasin et al. [14- 19]).

2. Coefficient inequality

In this section, we find the coefficient inequality for the class $P_{p,q}^{\zeta,k}(\beta, \delta, \lambda, \mu, b)$.

Theorem 2.1. Let the function $\varphi(z)$ be defined by (1.10). Then $\varphi(z) \in P_{p,q}^{\zeta,k}(\beta, \delta, \lambda, \mu, b)$ if and only if

$$\sum_{h=2}^{\infty} [h]_{p,q}^k \left\{ h[h]_{p,q} (1 + \mu) - \mu b - 1 \right\} [1 + (h-1)\beta(\delta - \lambda)]^{\zeta} a_h \leq \mu(1-b) \quad (2.1)$$

The result is sharp.

$$f(z) = z - \frac{\mu(1-b)}{[h]_{p,q}^k \left\{ h[h]_{p,q} (1 + \mu) - \mu b - 1 \right\} [1 + (h-1)\beta(\delta - \lambda)]^{\zeta}} z^h \quad (2.2)$$

Proof. Suppose that the inequality (2.1) holds. Then we have for $z \in \Theta$ and $|z| < 1$:

$$\begin{aligned} & \left| (1 - \beta(\delta - \lambda))z \left(\Lambda_{\beta, \delta, \lambda, p, q}^{\zeta, k+1} \varphi(z) \right)' + \beta(\delta - \lambda)z \left(\Lambda_{\beta, \delta, \lambda, p, q}^{\zeta, k+1} \varphi(z) \right)' - \right. \\ & \left. (1 - \beta(\delta - \lambda))\Lambda_{\beta, \delta, \lambda, p, q}^{\zeta, k} \varphi(z) - \beta(\delta - \lambda)\Lambda_{\beta, \delta, \lambda, p, q}^{\zeta, k} \varphi(z) \right| - \\ & \mu \left| (1 - \beta(\delta - \lambda))z \left(\Lambda_{\beta, \delta, \lambda, p, q}^{\zeta, k+1} \varphi(z) \right)' + \beta(\delta - \lambda)z \left(\Lambda_{\beta, \delta, \lambda, p, q}^{\zeta, k+1} \varphi(z) \right)' - \right. \\ & \left. b(1 - \beta(\delta - \lambda))\Lambda_{\beta, \delta, \lambda, p, q}^{\zeta, k} \varphi(z) - b\beta(\delta - \lambda)\Lambda_{\beta, \delta, \lambda, p, q}^{\zeta, k} \varphi(z) \right| \\ & = \left| \sum_{h=2}^{\infty} [h]_{p,q}^k (h[h]_{p,q} - 1) [1 + (h-1)\beta(\delta - \lambda)]^{\zeta} a_h z^h \right| - \\ & \mu \left| z(1-b) - \sum_{v=2}^{\infty} [h]_{p,q}^k (h[h]_{p,q} - b) [1 + (h-1)\beta(\delta - \lambda)]^{\zeta} a_h z^h \right| \\ & \leq \sum_{h=2}^{\infty} [h]_{p,q}^k \left\{ (1 + \mu)h[h]_{p,q} - 1 - b\mu \right\} [1 + (h-1)\beta(\delta - \lambda)]^{\zeta} a_h z^h - \mu(1-b) \\ & \leq 0 \end{aligned}$$

where $\Lambda_{\beta, \delta, \lambda, p, q}^{\zeta, k} \varphi(z)$ is given by (1.8).

This implies

$$\sum_{h=2}^{\infty} [h]_{p,q}^k \left\{ (1 + \mu)h[h]_{p,q} - 1 - b\mu \right\} [1 + (h-1)\beta(\delta - \lambda)]^{\zeta} a_h z^h \leq \mu(1-b)$$

which shows that $\varphi(z) \in P_{p,q}^{\zeta,k}(\beta, \delta, \lambda, \mu, b)$. For the converse, assume that

$$\begin{aligned}
& \left| \frac{(1-\beta(\delta-\lambda))z(\Lambda_{\beta,\delta,\lambda,p,q}^{\zeta,k+1}\varphi(z))' + \beta(\delta-\lambda)z(\Lambda_{\beta,\delta,\lambda,p,q}^{\zeta,k}\varphi(z))'}{(1-\beta(\delta-\lambda))\Lambda_{\beta,\delta,\lambda,p,q}^{\zeta,k}\varphi(z) + \beta(\delta-\lambda)\Lambda_{\beta,\delta,\lambda,p,q}^{\zeta,k}\varphi(z)} - 1 \right| \\
& \mu \left| \frac{(1-\beta(\delta-\lambda))z(\Lambda_{\beta,\delta,\lambda,p,q}^{\zeta,k+1}\varphi(z))' + \beta(\delta-\lambda)z(\Lambda_{\beta,\delta,\lambda,p,q}^{\zeta,k}\varphi(z))'}{(1-\beta(\delta-\lambda))\Lambda_{\beta,\delta,\lambda,p,q}^{\zeta,k}\varphi(z) + \beta(\delta-\lambda)\Lambda_{\beta,\delta,\lambda,p,q}^{\zeta,k}\varphi(z)} - b \right| \\
& \frac{\left| -\sum_{h=2}^{\infty} [h]_{p,q}^k (h[h]_{p,q} - 1) [1 + (h-1)\beta(\delta-\lambda)]^{\zeta} a_h z^h \right|}{\mu \left| z(1-b) - \sum_{h=2}^{\infty} [h]_{p,q}^k (h[h]_{p,q} - b) [1 + (h-1)\beta(\delta-\lambda)]^{\zeta} a_h z^h \right|} \\
& < 1
\end{aligned} \tag{2.3}$$

Since the $\operatorname{Re}(z) \leq |z|$ for all z , it follows from (2.3) that

$$\operatorname{Re} \left\{ \frac{\sum_{h=2}^{\infty} [h]_{p,q}^k (h[h]_{p,q} - 1) [1 + (h-1)\beta(\delta-\lambda)]^{\zeta} a_h z^h}{z(1-b) - \sum_{h=2}^{\infty} [h]_{p,q}^k (h[h]_{p,q} - b) [1 + (h-1)\beta(\delta-\lambda)]^{\zeta} a_h z^h} \right\} < 1. \tag{2.4}$$

By choosing values of z on the real axis and letting $|z| \rightarrow 1^-$ through the real values, we obtain

$$\begin{aligned}
& \sum_{h=2}^{\infty} [h]_{p,q}^k (h[h]_{p,q} - 1) [1 + (h-1)\beta(\delta-\lambda)]^{\zeta} a_h \\
& \leq (1-b)\mu - \mu \sum_{h=2}^{\infty} [h]_{p,q}^k (h[h]_{p,q} - b) [1 + (h-1)\beta(\delta-\lambda)]^{\zeta} a_h
\end{aligned}$$

This gives the required condition.

Corollary 2.2. Let the function $\varphi(z)$, defined by (1.10), be in the class $P_{p,q}^{\zeta,k}(\beta, \delta, \lambda, \mu, b)$. Then

$$a_h \leq \frac{\mu(1-b)}{[h]_{p,q}^k \left\{ h[h]_{p,q} (1+\mu) - \mu b - 1 \right\} [1 + (h-1)\beta(\delta-\lambda)]^{\zeta}}, \quad (h \geq 2). \tag{2.5}$$

The inequality in (2.1) is obtained for the function $\varphi(z)$ given by (2.2).

3. Growth and Distortion Theorems

Theorem 3.1. Let the function $\varphi(z)$, defined by (1.10), be in the class $P_{p,q}^{\zeta,k}(\beta, \delta, \lambda, \mu, b)$. Then for

$$|z| = r < 1,$$

$$\left| \Lambda_{\beta,\delta,\lambda,p,q}^{i,j} \varphi(z) \right| \geq r - \frac{\mu(1-b)}{[2]_{p,q}^{k-j} \left\{ 2[2]_{p,q} (1+\mu) - \mu b - 1 \right\} [1 + \beta(\delta-\lambda)]^{\zeta-i}} r^2 \tag{3.1}$$

and

$$\left| \Lambda_{\beta,\delta,\lambda,p,q}^{i,j} \varphi(z) \right| \leq r + \frac{\mu(1-b)}{[2]_{p,q}^{k-j} \left\{ 2[2]_{p,q} (1+\mu) - \mu b - 1 \right\} [1 + \beta(\delta-\lambda)]^{\zeta-i}} r^2, \tag{3.2}$$

$$(0 \leq i \leq \zeta, 0 \leq j \leq k, z \in \Theta)$$

The inequalities in (3.1) and (3.2) are obtained for $\varphi(z)$ given by

$$\varphi(z) = z - \frac{\mu(1-b)}{[2]_{p,q}^k \left\{ 2[2]_{p,q} (1+\mu) - \mu b - 1 \right\} [1 + (\beta(\delta-\lambda))]^{\zeta}} \tag{3.3}$$

Proof. Note that the function $\varphi(z) \in P_{p,q}^{\zeta,k}(\beta, \delta, \lambda, \mu, b)$ if and only if

$$\Lambda_{\beta,\delta,\lambda,p,q}^{i,j} \varphi(z) \in P_{p,q}^{\zeta,k}(\beta, \delta, \lambda, \mu, b)$$

and that

$$\Lambda_{\beta,\delta,\lambda,p,q}^{i,j} \varphi(z) = z - \sum_{h=2}^{\infty} [h]_{p,q}^j [1 + (h-1)\beta(\delta-\lambda)]^i a_h z^h \tag{3.4}$$

By the theorem 2.1

$$\begin{aligned} & [2]_{p,q}^k \left\{ 2[2]_{p,q} (1 + \mu) - \mu b - 1 \right\} [1 + (\beta(\delta - \lambda))]^\zeta \sum_{h=2}^{\infty} [h]_{p,q}^j (1 + \beta(\delta - \lambda))^i a_h \\ & \leq \sum_{h=2}^{\infty} [h]_{p,q}^k \left\{ 2[2]_{p,q} (1 + \mu) - \mu b - 1 \right\} [1 + (h-1)\beta(\delta - \lambda)]^\zeta a_h \leq \mu(1-b) \end{aligned} \quad (3.5)$$

Which implies,

$$\sum_{h=2}^{\infty} [h]_{p,q}^j [1 + \beta(\delta - \lambda)]^i a_h z^h \leq \frac{\mu(1-b)}{[2]_{p,q}^k \left\{ 2[2]_{p,q} (1 + \mu) - \mu b - 1 \right\} [1 + (\beta(\delta - \lambda))]^{\zeta-i}} \quad (3.6)$$

The assertions (3.1) and (3.2) of Theorem 4.1 would now follow readily from (3.4) and (3.6). Finally, we note that the equalities (3.1) and (3.2) are achieved for the function $\varphi(z)$, defined by

$$\Lambda_{\beta, \delta, \lambda, p, q}^{i, j} \varphi(z) = z - \frac{\mu(1-b)}{[2]_{p,q}^k \left\{ 2[2]_{p,q} (1 + \mu) - \mu b - 1 \right\} [1 + (\beta(\delta - \lambda))]^{\zeta-i}} z^2 \quad (3.7)$$

Hence, the proof has been completed.

Taking $i = j = 0$ in Theorem 2.1, we obtain this corollary.

Corollary 3.2.

Let $\varphi(z)$, defined by (1.10), be in the class $P_{p,q}^{\zeta, k}(\beta, \delta, \lambda, \mu, b)$. Then, for $|z| = r < 1$,

$$|\varphi(z)| \geq r - \frac{\mu(1-b)}{[2]_{p,q}^k \left\{ 2[2]_{p,q} (1 + \mu) - \mu b - 1 \right\} [1 + \beta(\delta - \lambda)]^\zeta} r^2 \quad (3.8)$$

and

$$|\varphi(z)| \leq r + \frac{\mu(1-b)}{[2]_{p,q}^k \left\{ 2[2]_{p,q} (1 + \mu) - \mu b - 1 \right\} [1 + \beta(\delta - \lambda)]^\zeta} r^2 \quad (3.9)$$

The equalities in (3.8) and (3.9) are achieved for the function $\varphi(z)$ given by (4.3).

4. Inclusion properties

We begin this section by showing the following inclusion relation.

Theorem 4.1. Let the hypotheses of theorem th1 be satisfied. Then

$$P_{p,q}^{\zeta, k}(\beta_1, \delta, \lambda, \mu_1, b) \supseteq P_{p,q}^{\zeta, k}(\beta_2, \delta, \lambda, \mu, b)$$

$$P_{p,q}^{\zeta, k}(\beta, \delta_1, \lambda, \mu, b) \supseteq P_{p,q}^{\zeta, k}(\beta, \delta_1, \lambda, \mu, b)$$

$$P_{p,q}^{\zeta, k}(\beta, \delta, \lambda_1, \mu, b) \supseteq P_{p,q}^{\zeta, k}(\beta, \delta, \lambda_2, \mu, b)$$

$$P_{p,q}^{\zeta, k}(\beta, \delta, \lambda, \mu_1, b) \supseteq P_{p,q}^{\zeta, k}(\beta, \delta, \lambda, \mu_2, b)$$

Proof. Let the function $\varphi(z)$, defined by (1.10), be in the class $P_{p,q}^{\zeta, k}(\beta, \delta, \lambda, \mu, b)$ and let $\beta_1 \geq \beta_2$.

Then, by theorem 2.1, we have

$$\begin{aligned} & \sum_{h=2}^{\infty} [h]_{p,q}^k \left\{ h[h]_{p,q} (1 + \mu) - \mu b - 1 \right\} [1 + (h-1)\beta_1(\delta - \lambda)]^\zeta a_h \\ & \leq \sum_{h=2}^{\infty} [h]_{p,q}^k \left\{ h[h]_{p,q} (1 + \mu) - \mu b - 1 \right\} [1 + (h-1)\beta_1(\delta - \lambda)]^\zeta a_h \\ & \leq \mu(1-b) \end{aligned}$$

Hence, $P_{p,q}^{\zeta, k}(\beta_1, \delta, \lambda, \mu_1, b) \supseteq P_{p,q}^{\zeta, k}(\beta_2, \delta, \lambda, \mu, b)$.

and

$$\begin{aligned} & \sum_{h=2}^{\infty} [h]_{p,q}^k \left\{ h[h]_{p,q} (1+\mu) - \mu b - 1 \right\} [1 + (h-1)\beta(\delta_1 - \lambda)]^{\zeta} a_h \\ & \leq \sum_{h=2}^{\infty} [h]_{p,q}^k \left\{ h[h]_{p,q} (1+\mu) - \mu b - 1 \right\} [1 + (h-1)\beta(\delta_2 - \lambda)]^{\zeta} a_h \\ & \leq \mu(1-b) \end{aligned}$$

Hence, $P_{p,q}^{\zeta,k}(\beta, \delta_1, \lambda, \mu, b) \supseteq P_{p,q}^{\zeta,k}(\beta, \delta_2, \lambda, \mu, b)$.

$$\begin{aligned} & \sum_{h=2}^{\infty} [h]_{p,q}^k \left\{ h[h]_{p,q} (1+\mu) - \mu b - 1 \right\} [1 + (h-1)\beta(\delta - \lambda_1)]^{\zeta} a_h \\ & \leq \sum_{h=2}^{\infty} [h]_{p,q}^k \left\{ h[h]_{p,q} (1+\mu) - \mu b - 1 \right\} [1 + (h-1)\beta(\delta - \lambda_1)]^{\zeta} a_h \\ & \leq 1 - \mu(1-b) \end{aligned}$$

Hence, $P_{p,q}^{\zeta,k}(\beta, \delta, \lambda_1, \mu, b) \supseteq P_{p,q}^{\zeta,k}(\beta, \delta, \lambda_2, \mu, b)$.

Employing a similar procedure, we can prove that $P_{p,q}^{\zeta,k}(\beta, \delta, \lambda, \mu_1, b) \supseteq P_{p,q}^{\zeta,k}(\beta, \delta, \lambda, \mu_2, b)$.

5. Closure Theorems

This section has begun with proving that the class $P_{p,q}^{\zeta,k}(\beta, \delta, \lambda, \mu, b)$ is closed under convex linear combinations.

Theorem 5.1. The class $P_{p,q}^{\zeta,k}(\beta, \delta, \lambda, \mu, b)$ is a convex set.

Proof. Let the functions

$$\varphi_{\varepsilon}(z) = z - \sum_{h=2}^{\infty} a_{\varepsilon,h} z^h \quad (a_{\varepsilon,h} \geq 0; \varepsilon = 1, 2; z \in \Theta) \quad (5.1)$$

be in the class $P_{p,q}^{\zeta,k}(\beta, \delta, \lambda, \mu, b)$. It is sufficient to show that the function $\gamma(z)$ defined by

$$\gamma(z) = \xi \varphi_1(z) + (1-\xi) \varphi_2(z) \quad (5.2)$$

is also in the class $P_{p,q}^{\zeta,k}(\beta, \delta, \lambda, \mu, b)$. Since, for $0 \leq \xi \leq 1$,

$$\gamma(z) = z - \sum_{h=2}^{\infty} \left\{ \xi a_{1,h} + (1-\xi) a_{2,h} \right\} z^h, \quad (5.3)$$

by using theorem 2.1, we have

$$\sum_{h=2}^{\infty} [h]_{p,q}^k \left\{ h[h]_{p,q} (1+\mu) - \mu b - 1 \right\} [1 + (h-1)\beta(\delta - \lambda)]^{\zeta} \left\{ \xi a_{1,h} + (1-\xi) a_{2,h} \right\} \leq \mu(1-b) \quad (5.4)$$

which means that $\gamma(z) \in P_{p,q}^{\zeta,k}(\beta, \delta, \lambda, \mu, b)$. Hence $P_{p,q}^{\zeta,k}(\beta, \delta, \lambda, \mu, b)$ is a convex set.

Theorem 5.2. Let $\varphi_1(z) = z$ and

$$\varphi_h(z) = z - \frac{\mu(1-b)}{[h]_{p,q}^k \left\{ h[h]_{p,q} (1+\mu) - \mu b - 1 \right\} [1 + (h-1)\beta(\delta - \lambda)]^{\zeta}} z^h, \quad (h \geq 2; k, \zeta \in \mathbb{N}_0) \quad (5.5)$$

for $0 \leq \mu < 1$ and $0 \leq \beta(\delta - \lambda) \leq 1$. Then $\varphi(z)$ is in the class $P_{p,q}^{\zeta,k}(\beta, \delta, \lambda, \mu, b)$ if and only if it could be expressed in the form:

$$\varphi(z) = \sum_{h=1}^{\infty} \omega_h \varphi_h(z), \quad (5.6)$$

where

$$\omega_h \geq 0 \quad (h \geq 1) \quad \text{and} \quad \sum_{h=1}^{\infty} \omega_h = 1 \quad (5.7)$$

Proof. Assume that

$$\begin{aligned}\varphi(z) &= \sum_{h=1}^{\infty} \omega_h \varphi_h(z) \\ &= z - \sum_{h=2}^{\infty} \frac{\mu(1-b)}{[h]_{p,q}^k \{h[h]_{p,q}(1+\mu) - \mu b - 1\} [1+(h-1)\beta(\delta-\lambda)]^\zeta} \omega_h z^h\end{aligned}$$

Then it follows that

$$\begin{aligned}\sum_{h=2}^{\infty} \frac{[h]_{p,q}^k \{h[h]_{p,q}(1+\mu) - \mu b - 1\} [1+(h-1)\beta(\delta-\lambda)]^\zeta}{\mu(1-b)} \\ \frac{\mu(1-b)}{[h]_{p,q}^k \{h[h]_{p,q}(1+\mu) - \mu b - 1\} [1+(h-1)\beta(\delta-\lambda)]^\zeta} \omega_h = \sum_{h=2}^{\infty} \omega_h = 1 - \omega_1\end{aligned}$$

Thus, by Theorem 2.1, $\varphi(z) \in P_{p,q}^{\zeta,k}(\beta, \delta, \lambda, \mu, b)$.

Conversely, suppose that $\varphi(z)$, defined by (1.10), is in the class $P_{p,q}^{\zeta,k}(\beta, \delta, \lambda, \mu, b)$. Then

$$a_h \leq \frac{\mu(1-b)}{[h]_{p,q}^k \{h[h]_{p,q}(1+\mu) - \mu b - 1\} [1+(h-1)\beta(\delta-\lambda)]^\zeta}, \quad (h \geq 2; k, \zeta \in \mathbb{N}_0).$$

considering

$$\omega_h = \frac{[h]_{p,q}^k \{h[h]_{p,q}(1+\mu) - \mu b - 1\} [1+(h-1)\beta(\delta-\lambda)]^\zeta}{\mu(1-b)} a_h \quad (h \geq 2; k, \zeta \in \mathbb{N}_0)$$

and

$$\omega_1 = 1 - \sum_{h=2}^{\infty} \omega_h$$

It's observable that $\varphi(z)$ can be expressed in (5.6). Which completes the proof.

6. Radii of close-to-convexity, starlikeness, and convexity

In this section, we shall determine the radii of close-to-convexity, starlikeness, and convexity for the functions belonging to the class $P_{p,q}^{\zeta,k}(\beta, \delta, \lambda, \mu, b)$.

Theorem 6.1. Let the function $\varphi(z)$, defined by (1.10), be in the class $P_{p,q}^{\zeta,k}(\beta, \delta, \lambda, \mu, b)$. Then

$\varphi(z)$ is close-to-convex of order σ ($0 \leq \sigma < 1$) in $|z| < r_1$, where

$$r_1 = \inf \left\{ \frac{(1-\sigma)h^{-1} [h]_{p,q}^k \{h[h]_{p,q}(1+\mu) - \mu b - 1\} [1+(h-1)\beta(\delta-\lambda)]^\zeta}{\mu(1-b)} \right\}^{\frac{1}{(h-1)}}, \quad (h \geq 2) \quad (6.1)$$

The result is sharp, with the extremal function $\varphi(z)$ given by (2.2).

Proof. We need to show that

$$|\varphi'(z) - 1| \leq 1 - \sigma \quad \text{for } |z| < r_1$$

where r_1 is given by (6.1). Then we yield from definition (1.10)

$$|\varphi'(z) - 1| \leq \sum_{h=2}^{\infty} h a_h |z|^{h-1}.$$

Thus,

$$|\varphi'(z) - 1| \leq 1 - \sigma$$

if

$$\sum_{h=2}^{\infty} \left(\frac{h}{1-\sigma} \right) a_h |z|^{h-1} \leq 1 \quad (6.2)$$

But, by Theorem 6.1, (6.2) holds true if

$$\left(\frac{h}{1-\sigma} \right) |z|^{h-1} \leq \frac{[h]_{p,q}^k \{h[h]_{p,q}(1+\mu) - \mu b - 1\} [1 + (h-1)\beta(\delta - \lambda)]^\zeta}{\mu(1-b)}$$

that is, if

$$|z| \leq \left(\frac{(1-\sigma)h^{-1} [h]_{p,q}^k \{h[h]_{p,q}(1+\mu) - \mu b - 1\} [1 + (h-1)\beta(\delta - \lambda)]^\zeta}{\mu(1-b)} \right)^{\frac{1}{(h-1)}} \quad (h \geq 2) \quad (6.3)$$

Theorem 6.1 follows readily from (6.4).

Theorem 6.2. Let $\varphi(z)$, defined by (1.10), be in the class $P_{p,q}^{\zeta,k}(\beta, \delta, \lambda, \mu, b)$. Then $\varphi(z)$ is a starlike of order σ ($0 \leq \sigma < 1$) in $|z| < r_2$, where

$$r_2 = \inf \left\{ \frac{(1-\sigma)[h]_{p,q}^k \{h[h]_{p,q}(1+\mu) - \mu b - 1\} [1 + (h-1)\beta(\delta - \lambda)]^\zeta}{\mu(h-\sigma)(1-b)} \right\}^{\frac{1}{(h-1)}}, \quad (h \geq 2) \quad (6.4)$$

The result is sharp, with the extremal function $\varphi(z)$ given by (2.2).

Proof. We need to show that

$$\left| \frac{z \varphi'(z)}{\varphi(z)} - 1 \right| \leq 1 - \sigma \quad \text{for } |z| < r_2$$

where r_2 is given by (6.4). Indeed, definition (1.10) implies that

$$\left| \frac{z \varphi'(z)}{\varphi(z)} - 1 \right| \leq \frac{\sum_{h=2}^{\infty} (h-1) a_h |z|^{h-1}}{1 - \sum_{h=2}^{\infty} a_h |z|^{h-1}}$$

Thus,

$$\left| \frac{z \varphi'(z)}{\varphi(z)} - 1 \right| \leq 1 - \sigma$$

if

$$\sum_{h=2}^{\infty} \left(\frac{h-\sigma}{1-\sigma} \right) a_h |z|^{h-1} \leq 1 \quad (6.5)$$

But, by Theorem 2.1, (6.5) holds true if

$$\left(\frac{h-\sigma}{1-\sigma} \right) |z|^{h-1} \leq \frac{[h]_{p,q}^k \{h[h]_{p,q}(1+\mu) - \mu b - 1\} [1 + (h-1)\beta(\delta - \lambda)]^\zeta}{\mu(h-\sigma)(1-b)} \quad (6.6)$$

that is, if

$$|z| \leq \left(\frac{(1-\sigma)[h]_{p,q}^k \{h[h]_{p,q}(1+\mu) - \mu b - 1\} [1 + (h-1)\beta(\delta - \lambda)]^\zeta}{\mu(h-\sigma)(1-b)} \right)^{\frac{1}{(h-1)}} \quad (h \geq 2) \quad (6.7)$$

Theorem 6.2 follows readily from (6.7).

Corollary 6.3. Let the function $\varphi(z)$, defined by (1.10), be in the class $P_{p,q}^{\zeta,k}(\beta, \delta, \lambda, \mu, b)$. Then $\varphi(z)$ is a convex of order σ ($0 \leq \sigma < 1$) in $|z| < r_2$, where

$$r_3 = \inf \left\{ \frac{(1-\sigma)h^{-1}[h]_{p,q}^k \{h[h]_{p,q}(1+\mu) - \mu b - 1\} [1 + (h-1)\beta(\delta - \lambda)]^\zeta}{\mu(h-\sigma)(1-b)} \right\}^{\frac{1}{(h-1)}}, (h \geq 2) \quad (6.8)$$

The result is sharp, with the extremal function $\varphi(z)$ given by (2.2).

7. Integral means inequality

For any two functions, φ and Γ , analytic in Θ , φ is said to be subordinate to Γ in Θ , written as $\varphi(z) \prec \Gamma(z)$, if there exists a Schwarz function $\omega(z)$, analytic in Θ , with

$\omega(0) = 0$ and $|\omega(z)| < 1$ for all $z \in \Theta$,

such that $\varphi(z) = \Gamma(\omega(z))$ for all $z \in \Theta$. Furthermore, if the function Γ is univalent in Θ , then we have the following equivalence [10]:

$\varphi(z) \prec \gamma(z) \Leftrightarrow \varphi(0) = \gamma(0)$ and $\varphi(\Theta) \subset \gamma(\Theta)$.

To prove the integral means inequality for functions belonging to the class $P_{p,q}^{\zeta,k}(\beta, \delta, \lambda, \mu, b)$, we need the following subordination result found by Littlewood [16].

Lemma 7.1. If the functions φ and Γ are analytic in Θ with $\varphi(z) \prec \Gamma(z)$, then for $\eta > 0$ and $z = re^{i\theta}$ ($0 < r < 1$),

$$\int_0^{2\pi} |\varphi(z)|^\eta d\theta \leq \int_0^{2\pi} |\gamma(z)|^\eta d\theta \quad (7.1)$$

By applying Theorem 2.1 with the extremal function and Lemma 7.1, we achieve the following theorem.

Theorem 7.2. Let $\{[h]_{p,q}^k \{h[h]_{p,q}(1+\mu) - \mu b - 1\} [1 + (h-1)\beta(\delta - \lambda)]^\zeta\}_{h=2}^\infty$ be a non-decreasing sequence. If $\varphi(z) \in P_{p,q}^{\zeta,k}(\beta, \delta, \lambda, \mu, b)$, then

$$\int_0^{2\pi} |\varphi(re^{i\theta})|^\eta d\theta \leq \int_0^{2\pi} |\gamma(re^{i\theta})|^\eta d\theta \quad (0 < r < 1; \eta > 0), \quad (7.2)$$

where

$$\varphi_*(z) = z - \frac{\mu(1-b)}{[2]_{p,q}^k \{2[2]_{p,q}(1+\mu) - \mu b - 1\} [1 + (\beta(\delta - \lambda))]^\zeta} z^2 \quad (7.3)$$

Proof. Let the function $\varphi(z)$, defined by (1.10), be in the class $P_{p,q}^{\zeta,k}(\beta, \delta, \lambda, \mu, b)$. Then we need to show that

$$\int_0^{2\pi} \left| 1 - \sum_{h=2}^\infty a_{h,h} z^{h-1} \right|^\eta d\theta \leq \int_0^{2\pi} \left| 1 - \frac{\mu(1-b)}{[2]_{p,q}^k \{2[2]_{p,q}(1+\mu) - \mu b - 1\} [1 + (\beta(\delta - \lambda))]^\zeta} z \right|^\eta d\theta \quad (7.4)$$

Thus, by applying Lemma 7.1, it would suffice to show that

$$1 - \sum_{h=2}^\infty a_{h,h} z^{h-1} \prec 1 - \frac{\mu(1-b)}{[2]_{p,q}^k \{2[2]_{p,q}(1+\mu) - \mu b - 1\} [1 + (\beta(\delta - \lambda))]^\zeta} z \quad (7.5)$$

If the subordination (7.5) holds true, then there exists an analytic function ω with $\omega(0) = 0$ and $|\omega(z)| < 1$ such that

$$1 - \sum_{h=2}^\infty a_{h,h} z^{h-1} = 1 - \frac{\mu(1-b)}{[2]_{p,q}^k \{2[2]_{p,q}(1+\mu) - \mu b - 1\} [1 + (\beta(\delta - \lambda))]^\zeta} \omega(z). \quad (7.6)$$

Using Theorem 2.1, we have

$$|\omega(z)| = \left| \sum_{h=2}^{\infty} \frac{[2]_{p,q}^k \{2[2]_{p,q}(1+\mu) - \mu b - 1\} [1 + (\beta(\delta - \lambda))]^{\zeta}}{\mu(1-b)} a_h z^{h-1} \right|$$

$$\leq |z| \sum_{h=2}^{\infty} \frac{[2]_{p,q}^k \{2[2]_{p,q}(1+\mu) - \mu b - 1\} [1 + (\beta(\delta - \lambda))]^{\zeta}}{\mu(1-b)} a_h \leq |z| < 1,$$

which proves the subordination (7.5). So the proof is completed.

References

1. Kanas, S. and Răducanu, D. **2014**. Some class of analytic functions related to conic domains. *Mathematica slovacica*, **64**(5): 1183-1196.
2. Jackson, F. H. **1909**. XI---on q-functions and a certain difference operator. *Earth and Environmental Science Transactions of the Royal Society of Edinburgh*, **46**(2): 253-281.
3. Aral, A., Gupta, V. and Agarwal, R. P. **2013**. *Applications of q-calculus in operator theory* (p. 262). New York: Springer.
4. Altinkaya, S. and Kanas, S. **2018**. Subclass of k-uniformly starlike functions defined by symmetric q-derivative operator. *Ukrains'kyi Matematychnyi Zhurnal*, **70**(11): 1499-1510
5. Yalçın, S., Vijaya, K. and Murugusundaramoorthy, G. **2018**. Certain class of analytic functions involving Salagean type q-difference operator. *Konuralp Journal of Mathematics*, **6**(2): 264-271.
6. Salagean, G. S. **1983**. Subclasses of univalent functions. In *Complex Analysis—Fifth Romanian-Finnish Seminar* (pp. 362-372). Springer, Berlin, Heidelberg.
7. Frasin, B.A. and Murugusundaramoorthy, G. **2000**. A subordination results for a class of analytic functions defined by q-differential operator, Submitted.
8. Yousef, F., Al-Shible, A. and Yalcin, S. **2019**. An Application of Jackson's (p, q)-Derivative to a Subclass of Starlike Functions with Negative Coefficients. *arXiv preprint arXiv: 2599556*.
9. Darus, M. and Ibrahim, R. W. **2009**. On subclasses for generalized operators of complex order. *Far East Journal of Mathematical Sciences*, **33**(3): 299-308.
10. Al-Oboudi, F. M. **2004**. On univalent functions defined by a generalized Sălăgean operator. *International Journal of Mathematics and Mathematical Sciences*, **2004**(27): 1429-1436.
11. Al-Hawary, T., Frasin, B. A. and Yousef, F. **2018**. Coefficients estimates for certain classes of analytic functions of complex order. *Afrika Matematika*, **29**(7-8): 1265-1271.
12. Al-Hawary, T., Yousef, F. and Frasin, B. A. **2018**. Subclasses of Analytic Functions of Complex Order Involving Jackson's (p, q)-derivative. *Available at SSRN* 3289803.
13. Aouf, M. K. and Srivastava, H. M. **1996**. Some families of starlike functions with negative coefficients. *Journal of mathematical analysis and applications*, **203**(3): 762-790.
14. Frasin, B. A., Al-Hawary, T. and Yousef, F. **2018**. Necessary and sufficient conditions for hypergeometric functions to be in a subclass of analytic functions. *Afrika Matematika*, 1-8.
15. Yousef, F., Amourah, A. A. and Darus, M. **2016**. Differential sandwich theorems for p-valent functions associated with a certain generalized differential operator and integral operator. *Italian Journal of Pure and Applied Mathematics*, **36**: 543-556.
16. Littlewood, J. E. **1925**. On inequalities in the theory of functions. *Proceedings of the London Mathematical Society*, **2**(1): 481-519.
17. Goodman, A. W. **1957**. Univalent functions and nonanalytic curves. *Proceedings of the American Mathematical Society*, **8**(3): 598-601.
18. Ruscheweyh, S. **1981**. Neighborhoods of univalent functions. *Proceedings of the American Mathematical Society*, **81**(4): 521-527.
19. Ahuja, O. P., Çetinkaya, A. and Ravichandran, V. **2018**. Harmonic Univalent Functions Defined by Post Quantum Calculus Operators. *arXiv preprint arXiv:1809.00298*.